






























DOSING FOR TYLENOL, IBUPROFEN & BENADRYL

ACETAMINOPHEN is the main ingredient in Tylenol. It is used as a pain reliever and fever reducer. It should **NOT** be used in children under 2 months of age. If your child requires acetaminophen for longer than three days, please contact the office. **The following can be used every 4-6 hours as needed:**

Weight (Lbs)	Infants' Tylenol (160 mg / 5 mL)	Children's Tylenol (160 mg / 5 mL)	Tylenol Chewables 160 mg	Tylenol Dissolve Packs 160 mg
6-11 lbs	1.25 mL 	DO NOT USE	DO NOT USE	DO NOT USE
12-17 lbs	2.5 mL 	DO NOT USE	DO NOT USE	DO NOT USE
18-23 lbs	3.75 mL 	DO NOT USE	DO NOT USE	DO NOT USE
24-35 lbs	5 mL 	5 mL 	1 tablet 	DO NOT USE
36-47 lbs	DO NOT USE	7.5 mL 	1 1/2 tablets 	DO NOT USE
48-59 lbs	DO NOT USE	10 mL 	2 tablets 	2 powders 
60-71 lbs	DO NOT USE	12.5 mL 	2 1/2 tablets 	2 powders 
72-95 lbs	DO NOT USE	15 mL 	3 tablets 	3 powders 

DO NOT GIVE MORE THAN 5 DOSES IN 24 HOURS

IBUPROFEN is the active ingredient in Advil and Motrin. Ibuprofen is used to decrease pain and swelling and to reduce fevers. It should not be used in children less than 6 months old without discussing first with your practitioner. **The following can be used every 6-8 hours as needed:**

Weight (lbs)	Infants' Ibuprofen (50 mg / 1.25 mL)	Children's Ibuprofen (100 mg / 5 mL)	Ibuprofen Chewables (100 mg tablets)
12-17 lbs	1.25 mL 	DO NOT USE	DO NOT USE
18-23 lbs	1.875 mL 	DO NOT USE	DO NOT USE
24-35 lbs	DO NOT USE	5 mL 	1 tablet 
36-47 lbs	DO NOT USE	7.5 mL 	1 1/2 tablets 
48-59 lbs	DO NOT USE	10 mL 	2 tablets 
60-71 lbs	DO NOT USE	12.5 mL 	2 1/2 tablets 
72-95 lbs	DO NOT USE	15 mL 	3 tablets 

DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS

BENADRYL (diphenhydramine) is used for hives and to reduce the itchiness associated with certain rashes. **Benadryl can be given every 6-8 hours as needed.** Benadryl can cause drowsiness, dizziness, or paradoxical excitement.

Weight (lbs)	Benadryl Liquid (12.5 mg / tsp)	Benadryl Chewable (12.5 mg)	Benadryl Tabs (25 mg)
12-17 lbs	1/2 tsp	DO NOT USE	DO NOT USE
18-23 lbs	3/4 tsp	DO NOT USE	DO NOT USE
24-35 lbs	1 tsp	1 tablet	DO NOT USE
36-47 lbs	1 1/2 tsp	1 1/2 tablets	2 tablets
48-59 lbs	2 tsp	2 tablets	2 1/2 tablets

VISIT OUR WEBSITE:

WWW.DESERTPEDIATRICSLV.COM